

AN ABS ALGORITHM FOR A CLASS OF SYSTEMS OF STOCHASTIC LINEAR EQUATIONS

Hai-Shan Han¹, Antonino Del Popolo² and Zun-Quan Xia³

Abstract. This paper is to explore a model of the ABS Algorithms dealing with the solution of a class of systems of linear stochastic equations $A\xi = \eta$ when η is a m -dimensional normal distribution. It is shown that the stepsize α_i is distributed as $N(u_i, \sigma_i)$ (being u_i the expected value of α_i and σ_i its variance) and the approximation to the solutions ξ_i is distributed as $N_n(U_i, \Sigma_i)$ (being U_i the expected value of ξ_i and Σ_i its variance) , for this algorithm model.

Key words: ABS algorithm, stochastic linearly system of equations, distribution, probability.

AMS Subject Classification (2000): 60H35 65H10 65F10.

1 Introduction

Since the early fifties, the theory of stochastic equations, and in particular stochastic integral equations, has been the stage for intense activity. This has been stimulated in part by an immense wealth of applications beginning with the Langevin equation in statistical mechanics and Wiener's emphasis on the role of randomness in the problems of cybernetics. Applications of random equations are now found in many areas of engineering, physical and biological sciences and systems theory.

Historically the study of random equations (stochastic integral equations, stochastic operator equations, stochastic difference equations, etc.) started with that of integral equations. The theory of the last has grown along two parallel branches. In one,

¹CORA, Department of Applied Mathematics, Dalian University of Technology Dalian 116024, China; Department of Mathematics, Inner Mongolia University of Nationalities, Tongliao 028000 China

²Dept. of Mathematics, University of Bergamo, via dei Caniana 2, 24127 Bergamo, Italy;
Istanbul Technical University, Ayazaga Campus, Faculty of Science and Letters, 34469 Maslak/Istanbul, Turkey

³CORA, Department of Applied Mathematics, Dalian University of Technology Dalian 116024, China, zqxiazhh@dlut.edu.cn

initiated by K. Ito in 1951, the source of randomness is a white noise term leading to an important class of stochastic integral equations which has developed hand in hand with the theory of Markov processes. In the other, classical linear and nonlinear integral equations with random right-hand sides, random kernels or defined on random domains are studied.

The application of stochastic equations is more extensive than deterministic equations, because random factors are usually included in many natural phenomena.

The study of random linear equations can at least be traced back to the works of [Bellman 54], [Rosenbloom 55], [Kalman 62]. Random linear difference equations arises naturally in many areas of engineering sciences including automatic control and signal processing, and have drawn much attention from researchers in the past several decades [Brandt 86].

Attempts were made by many scientists and mathematicians to develop and unify the theory of random equations employing concepts and methods of probability theory and functional analysis. The Prague School of probabilists under Spacek initiated a systematic study using probabilistic operator equations as models for various systems. This development was further energized by the survey article by [BhRe 72] on various treatments of random equations under the framework of functional analysis.

Several techniques have been proposed to solve stochastic linear equations (see for instance [Brandt 1986], [Verma 1996]). Stability problems of random linear equations have also been studied in [Guo 90], [Guo 93], [GuoLj 95].

An approach that has never been tried to solve stochastic linear equations is through the use of ABS methods, introduced by Abaffy, Broyden and Spedicato in 1984 in the form of a large class of methods for solving linear determined or underdetermined systems. The original formulation of 1984 has been later generalized giving a larger class of methods that essentially contains all possible methods building a sequence of approximations x_i to the solution having the following property: for x_1 arbitrary, the vector x_{m+1} solves the given system. ABS methods have been applied the least squares problem, to non-linear equations, to linear and non-linear programming and eigenvalue problems.

Before showing how the ABS algorithm can be used to solve a system of stochastic

equations of the form

$$A\xi(w) = \eta(w) \quad (1.1)$$

where $\xi(w)$, $\eta(w)$ are random vectors, we recall the ABS algorithm.

Consider the general linear system,

$$Ax = b \quad \text{or} \quad a_i^T x = b_i, \quad i = 1, \dots, m \quad (1.2)$$

where $x \in \mathbb{R}^m$, $b \in \mathbb{R}^n$, $m \leq n$ and $A \in \mathbb{R}^{m \times n}$, $A = (a_1, a_2, \dots, a_m)^T$.

The class of ABS algorithms to solve Eq. (1.2) was originally introduced by Abaffy, Broyden and Spedicato (1984), see [AbBs84] and [AbSp89]. The iterate scheme of the basic ABS class of algorithm is defined as follows:

Basic (unscaled) ABS Class of Algorithms: [AbBs84], [AbBs89]:

(A) Initialization.

Give an arbitrary vector $x_1 \in \mathbb{R}^n$, and an arbitrarily nonsingular matrix $H_1 \in \mathbb{R}^{n,n}$. Set $i = 1$.

(B) Compute $s_i = H_i a_i$ and $\tau_i = \tau^T e_i = a_i^T x_i - b^T e_i$

(C) Check the compitability of the system of linear equations.

If $s_i \neq 0$ then goto (D).

If $s_i = 0$ and $\tau_i = 0$ then set $x_{i+1} = x_i$, $H_{i+1} = H_i$ and goto (F), the i -th equation is a linear combination of the previous equations. Otherwise stop, the system has no solution.

(D) Compute the search vector $p_i \in \mathbb{R}^n$ by $p_i = H_i^T z_i$ where z_i , the parameter of Broyden, is arbitrary satisfying the condition $z_i^T H_i a_i \neq 0$

(E) Update the approximation of the solution x_i by $x_{i+1} = x_i - \alpha_i p_i$, where the stepsize α_i is computed by $\alpha_i = \tau_i / a_i^T p_i$ If $i = m$ stop; x_{m+1} solves the system.

(F) Update the (Abaffian) matrix H_i . Compute $H_{i+1} = H_i - H_i a_i w_i^T H_i / w_i^T H_i a_i$ where $w_i \in \mathbb{R}^n$, the parameter of Abaffy, is arbitrary satisfying the condition

$$w_i^T H_i a_i = 1 \text{ or } \neq 0$$

(G) Increment the index i by one and goto (B).

We define n by i matrices A_i , W_i and P_i by

$$A_i = (a_1, \dots, a_i)^T, \quad W_i = (w_1, \dots, w_i), \quad P_i = (p_1, \dots, p_i)$$

Some properties of the above recursion (see for instance, Abaffy and Spedicato (1989), [AbSp 89]), that are the basic formulae to be used later on, are listed below .

a. Implicit factorization property $A_i^T P_i = L_i$ with L_i nonsingular lower triangular.

b. Null space characterizations

$$\mathcal{N}(H_{i+1}) = \mathcal{R}(A_i^T), \quad \mathcal{N}(H_{i+1}^T) = \mathcal{R}(W_i),$$

$$\mathcal{N}(A_i) = \mathcal{R}(H_{i+1}^T)$$

where $\mathcal{N}$ = Null and $\mathcal{R}$ =Range.

c. The linear variety containing all solutions to $Ax = b$ consists of the vectors of the form $x = x_{i+1} + H_{i+1}^T q$ where $q \in \mathbb{R}^n$ is arbitrary.

As we shall show in the rest of the paper, this method can be used to solve a linear stochastic system in the following way. For a linear system

$$A\xi = \eta$$

with a $m \times n$ matrix of arbitrary rank with $m \leq n$, the so-called ABS algorithm produces a solution

$$\xi = A^I \eta,$$

with A^I some generalised inverse. Now it is quite clear that if η is a normally distributed vector, so is ξ , or in other words in the original system there are also normally distributed vector so is ξ as solutions. The mean of ξ can obviously be computed from the mean of η in the usual deterministic way (save for a component in the kernel of A). So the only interesting question left is how to compute the covariance of ξ from that of η . Formally it is $A^I V (A^I)^T$ if the covariance of η is V . The paper gives a recursive formula through the ABS algorithm for the special case that the components of η are uncorrelated, i. e. $V = I$.

This paper is organized as follows. In Section 2, some basic definitions and operations that will be used below are presented. A basic ABS algorithm, called ABS-S for stochastic linear equations is given in Section 3. Some properties of the approximations to solutions and α_i , are given in section 4. In section 5, the ABS-S algorithm is illustrated by an example. In section 6, α_i is discussed and same results are also given.

2 Preliminaries

Consider a class of systems of stochastic linear equations being of the form

$$A\xi(w) = \eta(w) \quad (2.1)$$

In this paper, we present an ABS algorithm under the condition that $\eta(w)$ has a m -dimensional normal distribution. It will be shown that the iteration step length $\alpha_i \sim N(u_i, \sigma_i)$ (being u_i the expected value of α_i and σ_i its variance) and the iteration solution ξ_i is $N_n(U_i, \Sigma_i)$, $i=1,2,\dots,m+1$.

We show that (u_i, σ_i) and (U_i, Σ_i) are determined by four iterative formulae, that ξ_{i+1} is a solution of the first i equations and we discuss the step length α_i .

Definition 2.1[GoMi84] Let $(\Omega, \mathcal{A}, \mathcal{P})$ be a probability space and let $\xi: \Omega \rightarrow R$ be a real-valued function on Ω . We say that ξ is a random variable on Ω iff for every $x \in R$, $\{\omega : \xi(\omega) \leq x\} \in \mathcal{A}$.

Definition 2.2 If a random variable ξ has a normal distribution with expectation μ and variance σ^2 , then it is denoted by $\xi \sim N(\mu, \sigma^2)$.

Definition 2.3[GoMi84] $\eta = (\eta_1, \eta_2, \dots, \eta_m)$ has a joint $N(\mu, \Sigma)$ density iff there exists an $m \times n$ matrix $A(m \leq n)$ of rank m such that $\eta = A\xi + \mu$ and $\xi = (\xi_1, \xi_2, \dots, \xi_n)$ is a vector of independent $N(0,1)$ random variables. Equivalently ξ is a $N(0, I_n)$ multiple normal density.

Definition 2.4 A random variable of random distribution ξ is said to be equal to the random variable of random distribution η , denoted by $\xi = \eta$, if the distributions of ξ and η are the same and $E\xi=E\eta$ and $D\xi=D\eta$, where E and D stands for expectation value and variance, respectively.

Definition 2.5 Consider a system of random linear equations $A\xi=\eta$. Suppose that η has a certain distribution. If the distribution of ξ , denoted by ξ_0 , is given, then ξ_0 is a solution of equations $A\xi=\eta$.

Property 2.1[GoMi84] If $\xi = \eta + a$, $a \in \mathcal{R}$, then $E\xi=E\eta+a$, $D\xi=D\eta$.

Property 2.2[Ande 84] If (ξ, η) is a 2-dimensional normal distribution, then $a\xi + b\eta$ is also a normal distribution, where a and b are constants.

3 ABS-S Algorithm to solve System of Stochastic Linear Equations

Consider a system of stochastic linear equations

$$A\xi = \eta \tag{3.1}$$

where $\eta = (\eta_1, \eta_2, \dots, \eta_m)^T$ is a m -dimensional stochastic vector and $A = (a_1, a_2, \dots, a_m)^T \in R^{m,n}$

The ABS-S algorithm, based on the basic ABS algorithm, is defined, as follows.

ABS-S Algorithm to solve systems of stochastic linear equations

(A1) Initialization.

Give an arbitrary stochastic vector $\xi_1 \in \mathbb{R}^n$, and an arbitrarily nonsingular matrix $H_1 \in R^{n,n}$. Set $i = 1$

(B1) Compute the two quantities:

$$s_i = H_i a_i, \tau_i = a_i^T \xi_i - \eta_i$$

(C1) Check the compatibility of the system of linear equations.

If $s_i \neq 0$ then goto (D).

If $s_i = 0$ and $Prob.(\tau_i = 0) = 1$ then set

$$\begin{aligned}\xi_{i+1} &= \xi_i \\ H_{i+1} &= H_i\end{aligned}$$

and go to (F), the i -th equation is a linear combination of the previous equations.

Otherwise stop, the system has no solution.

(D1) Compute the search vector $p_i \in \mathbb{R}^n$ by $p_i = H_i^T z_i$ where z_i , the parameter of Broyden, is arbitrary satisfying the condition

$$z_i^T H_i a_i \neq 0$$

(E1) Update the random approximation ξ_i of a solution by

$$\xi_{i+1} = \xi_i - \alpha_i p_i$$

where the stepsize α_i is computed by

$$\alpha_i = \frac{\tau_i}{a_i^T p_i}$$

If $i = m$ stop; ξ_{m+1} solves the system(3.1).

(F1) Update the (Abaffian) matrix H_i . Compute

$$H_{i+1} = H_i - \frac{H_i a_i^T w_i H_i}{w_i^T H_i a_i}$$

where $w_i \in \mathbb{R}^n$, the parameter of Abaffy, is arbitrary satisfying the condition

$$w_i^T H_i a_i = 1 \text{ or } \neq 0$$

(G1) Increment the index i by one and go to (B1).

4 The properties of α_i and ξ_i

Let $\eta \sim N_m(v, I_m)$ where $\eta = (\eta_1, \eta_2, \dots, \eta_m)^T$ and $v = (v_1, v_2, \dots, v_m)^T$

Proposition 4.1

$$\begin{aligned}\tau_1 &= a_1^T \xi_1 - \eta_1 \sim N(a_1^T \xi_1 - v_1, 1) \\ \alpha_1 &= \frac{\tau_1}{a_1^T p_1} \sim N\left(\frac{a_1^T \xi_1 - v_1}{a_1^T p_1}, \frac{1}{(a_1^T p_1)^2}\right)\end{aligned}$$

Proof. From $\eta \sim N_m(v, I_m)$, we know that $\eta_i \sim N(v_i, 1), i = 1, 2, \dots, m$. By property 2.2, τ_1 has a normal distribution and $\eta \sim N(v, I_m)$, $\eta_i \sim N(v_i, 1), i = 1, 2, \dots, m$

$$\begin{aligned}E\tau_1 &= E(a_1^T \xi_1 - \eta_1) = a_1^T \xi_1 - v_1; \\ D\tau_1 &= D(a_1^T \xi_1 - \eta_1) = 1; \\ u_1 = E\alpha_1 &= E((a_1^T p_1)^{-1} \tau_1) = (a_1^T p_1)^{-1} (a_1^T \xi_1 - v_1); \\ \sigma_1 = D\alpha_1 &= D((a_1^T p_1)^{-1} \tau_1) = (a_1^T p_1)^{-2}.\end{aligned}$$

Therefore

$$\begin{aligned}\alpha_1 &= (a_1^T p_1)^{-1} \tau_1 \sim N(u_1, \sigma_1) \\ \tau_1 &= a_1^T \xi_1 - \eta_1 \sim N(E\tau_1, D\tau_1)\end{aligned}$$

The proof is completed.

Proposition 4.2 If ξ_2 is generated by the ABS-S algorithm, then

$$\xi_2 \sim N_n\left(\xi_1 - \frac{a_1^T \xi_1 - v_1}{a_1^T p_1} p_1, \frac{p_1 p_1^T}{(a_1^T p_1)^2}\right)$$

Proof. By using Proposition 4.1, we have that $\alpha_1 \sim N(E\alpha_1, D\alpha_1)$. By property

2.2, since $\xi_2 = \xi_1 - \alpha_1 p_1$ has a normal distribution and

$$\begin{aligned}U_2 = E\xi_2 &= E(\xi_1 - \alpha_1 p_1) = E(\xi_1) - E(\alpha_1 p_1) = \xi_1 - (a_1^T p_1)^{-1} (a_1^T \xi_1 - v_1) p_1 \\ \Sigma_2 = D\xi_2 &= D(\xi_1 - \alpha_1 p_1) = (a_1^T p_1)^{-2} p_1 p_1^T\end{aligned}$$

it follows that

$$\xi_2 \sim N(U_2, \Sigma_2)$$

The proof is completed.

Proposition 4.3 $a_i p_j = 0, \quad i < j$

Proof. see [AbBs 84], [AbSp 89]

Proposition 4.4 If τ_i is generated by the ABS-S algorithm, then τ_i has a normal distribution, and

$$\tau_i \sim N(a_i^T \xi_1 - a_i^T \sum_{j=1}^{i-1} (E\alpha_j) p_j - v_i, 1 + a_i^T \sum_{j,k=1}^{i-1} \text{cov}(\alpha_j p_j, \alpha_k p_k) a_i), i \geq 2$$

where

$$\text{cov}(\alpha_j p_j, \alpha_k p_k) = \rho_{jk} p_j^T p_j p_k^T p_k D(\alpha_j) D(\alpha_k)$$

where ρ_{jk} is the correlation coefficient of $\alpha_j p_j$ and $\alpha_k p_k$

Proof. By the ABS-S algorithm, we have that $\tau_i = a_i^T \xi_i - \eta_i$, $\xi_{i+1} = \xi_i - \alpha_i p_i$ therefore

$$\tau_i = a_i^T \xi_i - \eta_i = a_i^T (\xi_{i-1} - \alpha_{i-1} p_{i-1}) - \eta_i = \dots = a_i^T (\xi_1 - \sum_{j=1}^{i-1} (\alpha_j p_j)) - \eta_i$$

Because $\alpha_j, j < i$ has a normal distribution, property 2.2, $\eta_i \sim N(v_i, 1)$ hence τ_i has a normal distribution, and

$$\begin{aligned} E\tau_i &= E(a_i^T (\xi_1 - \sum_{j=1}^{i-1} (\alpha_j p_j)) - \eta_i) = a_i^T (\xi_1 - \sum_{j=1}^{i-1} (E\alpha_j) p_j) - v_i \\ D\tau_i &= D(a_i^T (\xi_1 - \sum_{j=1}^{i-1} (\alpha_j p_j)) - \eta_i) = a_i^T [\sum_{j,k=1}^{i-1} \text{cov}(\alpha_j p_j, \alpha_k p_k)] a_i + 1. \end{aligned}$$

The proof is completed.

Proposition 4.5 If α_i is generated by the ABS-S algorithm, then

$$\alpha_i \sim N\left(\frac{a_i^T \xi_1 - a_i^T \sum_{j=1}^{i-1} (E\alpha_j) p_j - v_i}{a_i^T p_i}, \frac{a_i^T [\sum_{j,k=1}^{i-1} \text{cov}(\alpha_j p_j, \alpha_k p_k)] a_i + 1}{(a_i^T p_i)^2}\right), i \geq 2$$

Proof. Since $\alpha_i = (a_i^T p_i)^{-1} \tau_i$ and property 2.2, we have that the proposition holds.

The proof is completed.

Theorem 4.1 *If ξ_{i+1} is generated by the ABS-S algorithm, then*

$$\xi_{i+1} \sim N_n(\xi_1 - \sum_{j=1}^{j=i} (E\alpha_j)p_j, \sum_{j,k=1}^i \text{cov}(\alpha_j p_j, \alpha_k p_k))$$

or

$$\xi_{i+1} \sim N_n(E\xi_i - (E\alpha_i)p_i, D\xi_i + p_i D(\alpha_i)p_i^T - 2\text{cov}(\xi_i, \alpha_i p_i)), i \geq 2$$

Proof. Since $\xi_i = \xi_{i-1} - \alpha_{i-1}p_{i-1} = \xi_{i-2} - \alpha_{i-2}p_{i-2} - \alpha_{i-1}p_{i-1} = \dots = \xi_1 - \sum_{j=1}^{j=i-1} \alpha_j p_j$, and $\alpha_j, j = 1, 2, \dots, i-1$ have a normal distribution, we have that ξ_i has a normal distribution and

$$E\xi_i = E(\xi_1 - \sum_{j=1}^{j=i-1} (\alpha_j p_j)) = \xi_1 - \sum_{j=1}^{j=i-1} E(\alpha_j p_j)$$

$$D\xi_i = D(\xi_1 - \sum_{j=1}^{j=i-1} (\alpha_j p_j)) = \sum_{j,k=1}^{i-1} \text{cov}(\alpha_j p_j, \alpha_k p_k)$$

in terms of property 2.2. The proof is completed.

Theorem 4.2 *ξ_{i+1} is the solution of the first i equations*

Proof. Since $\xi_{i+1} = \xi_i - \alpha_i p_i$, one has that $a_i^T \xi_{i+1} = a_i^T (\xi_i - \alpha_i p_i) = \eta_i$. If $l < i$, then one has that $a_l^T \xi_{i+1} = a_l^T (\xi_i - \alpha_i p_i) = a_l^T \xi_i$. Therefore, $E(a_l^T \xi_{i+1}) = E(a_l^T \xi_i)$ and $D(a_l^T \xi_{i+1}) = D(a_l^T \xi_i)$, one has

$$E(a_l^T \xi_{i+1}) = E(a_l^T \xi_i) = \dots = E(a_l^T \xi_l) = v_l,$$

$$D(a_l^T \xi_{i+1}) = D(a_l^T \xi_i) = \dots = D(a_l^T \xi_l) = \dots = D(a_1^T \xi_1)$$

The proof is completed.

Proposition 4.6 The ABS-S algorithm terminates in finite steps.

Proof. The assertion comes directly from Definition (2.5), i.e. according to the definition of a solution of the system of the stochastic linear equations $A\xi = \eta$. The proof is completed.

5 Conclusion and Discussion

5.1 On the step length

Let

$$E\alpha_i = \frac{a_i^T \xi_1 - v_i}{a_i^T p_i}$$

$$D\alpha_i = \frac{1 + a_i^T \sum_{j,k=1}^{i-1} \text{cov}(\alpha_j p_j, \alpha_k p_k) a_i}{(a_i^T p_i)^2}$$

It follows that

$$\alpha_i \sim N(E\alpha_i, \sqrt{D\alpha_i})$$

and hence

$$P(E\alpha_i - \sqrt{D\alpha_i} \leq \alpha_i \leq E\alpha_i + \sqrt{D\alpha_i}) = 0.6827 \quad (5.1)$$

$$P(E\alpha_i - 2\sqrt{D\alpha_i} \leq \alpha_i \leq E\alpha_i + 2\sqrt{D\alpha_i}) = 0.9545 \quad (5.2)$$

$$P(E\alpha_i - 3\sqrt{D\alpha_i} \leq \alpha_i \leq E\alpha_i + 3\sqrt{D\alpha_i}) = 0.9973 \quad (5.3)$$

Therefore, we obtain that

$$\alpha_i \in [E\alpha_i - 3\sqrt{D\alpha_i}, E\alpha_i + 3\sqrt{D\alpha_i}]$$

in other words, α_i are determined by the initial vector ξ_1 , and the first component v_1 of η as well as $a_i^T p_i$.

5.2 Main results

1. the system of stochastic linear equations (1.1) with $\eta \sim N_m(v, I_m)$ can be solved by the ABS-S algorithm under some assumptions.
2. The solution ξ_i generated by the ABS-S algorithm has a normal distribution, if $\eta \sim N_m(v, I_m)$.
3. The step length α_i generated by the ABS-S algorithm has a normal distribution, if $\eta \sim N_m(v, I_m)$.
4. Since the matrix A is non-random, H_i, p_i generated by ABS-S algorithm are non-random, thus the ABS-S algorithm to solve $A\xi = \eta$ has some properties that are the same as in the case of the basic ABS algorithm.

5.3 Open problems

1. Suppose that η has a normal distribution, but the components of η are dependent. An open problem is whether the system $A\xi = \eta$ has a solution, and if it has whether it can be solved by the ABS-S algorithm.
2. Suppose that the components of η have different distributions. Another open problem is whether the system $A\xi = \eta$ has a solution, and if it has whether it can be solved by the ABS-S algorithm.
3. Let A be a random matrix, another open problem is whether the system $A\xi = \eta$ has a solution, and if it has whether it can be solved by the ABS-S algorithm.
4. Suppose η is a function of a random variable. Another open problem is whether the system $A\xi = \eta$ has a solution, and if it has whether it can be solved by the ABS-S algorithm.

References

- [AbBs 84] Abaffy, J., Broyden, C.G., and Spedicato, E.: A class of direct methods for linear systems, *Numerische Mathematik* 45, 361-376 (1984).
- [AbSp 89] Abaffy, J., and Spedicato, E.: *ABS Projection Algorithms: Mathematical Techniques for Linear and Nonlinear Algebraic Equations*, Ellis Horwood, Chichester (1989).
- [Bellman 54] ellman, R.: Limit theorems for non-commutative operators I, *Duke Math. J.* Vol. 21, pp. 491-500, (1954).
- [Brandt 86] Brandt, A.: The Stochastic Equation $Y_{n+1} = A_n Y_n + B_n$ with Stationary Coefficients, *Adv. Appl. Prob.* 18(2), 211-220 (1986)
- [BhRe 72] Bharucha-Reid, A.T.: *Random Integral Equations*, Academic Press, New York, (1972).
- [BhRe 79] Bharucha-Reid, A.T.: *Approximate Solution of Random Equations*, North-Holland (1979).

- [Char 91] Charles, M. Goldie: Implicit Renewal Theory and Tails of solutions of Random Equations, *Ann. Appl. Prob.* 23(1), 126-166, (1991).
- [DaBr 72] Brillinger, D.R., On the Number of Solutions of Systems of Random Equations, *Ann. Math. Stat.* 43(2), 534-540 (1972).
- [Guo 90] Guo, L.: Estimating time-varying parameters by Kalman filter based algorithms: Stability and convergence, *IEEE Trans. on Automatic Control*, Vol. 135, No.2, pp 141-147 (1990).
- [Guo 93] Guo, L.: Time-Varying Stochastic Systems: Stability, Estimation and Control (in Chinese), Jilin Science and Technology Press (1993).
- [Guo 94] Guo, L.: Stability of recursive stochastic tracking algorithms, *SIAM J. on Control and Optimization*, Vol. 32, 5, 1195-1225 (1994).
- [GuoLj 95] Guo, L., and Ljung, L.: Exponential stability of general tracking algorithms, *IEEE Trans. on Automatic Control*, vol 140 (1995).
- [Kalman 62] Kalman, R.E., Control of randomly varying linear dynamical systems, *Proc. Sympos. Appl. Math.*, Vol. 13, Amer. Math. Soc. Providence, R.I., 287-298 (1962).
- [KlPe 2004] Kluppelberg, C., and Pergamenchtchikov, S.: The Tail of the Stationary Distribution of a Random Coefficient AR(q) Model, *Ann. Appl. Probab.* 14(2), 971-1005 (2004).
- [KlPeRe 2003] Kluppelberg, C., and Pergamenchtchikov, S.: Renewal Theory for Functionals of A Markov Chain with Compact State Space, *Ann. Probab.* 31(4), 2270-2300, (2003).
- [Rosebloom 1955] Rosenbloom, A., et al.: Analysis of linear systems with randomly varying inputs and parameters, *IRE convention record*, Pt. 4, pp. 106.
- [Sped 93] Spedicato, E.: Ten years of ABS methods: a review of theoretical results and computational achievements, *Surveys on Mathematics for Industry*, 3, 217-232 (1993).

- [Sped 97] Spedicato, E.: ABS algorithms from Luoyang to Beijing, OMS 8, 87-97 (1997).
- [Sped 01] Spedicato, E.: Twenty years of ABS algorithms: A bibliography, QDMSIA 01/13, University of Bergamo (2001)
- [SpXZ 00] Spedicato, E., Xia, Z., and Zhang, L.: ABS algorithms for linear equations and optimization, J. of Computational and Applied Mathematics, 124, 155-170 (2000).
- [Chun 74] Chung, K.L.: A Course in Probability Theory, Second Edition, Academic press (1974).
- [Verma 96] Verma R.U.: Stochastic approximation-solvability of linear random equations involving numerical ranges, Journal of Applied Mathematics and Stochastic Analysis, 10:1, 47-55 (1996).
- [Wjd90] Wang, J.D.: Stochastic Programming(Chinese), Nan Jing University Press in China, (1990).
- [ZhFa 90] Zhang, Y.T., and Fang K.T.: Multivariate Statistical Analysis Inference(Chinese), Science Press in China (1999).